* * * *

NEWS 1 Web Page for STN Seminar Schedule - N. America

NEWS 2 JAN 12 Match STN Content and Features to Your Information

Needs, Quickly and Conveniently

NEWS 3 JAN 25 Annual Reload of MEDLINE database NEWS 4 FEB 16 STN Express Maintenance Release, Version 8.4.2, Is

Now Available for Download

NEWS 5 FEB 16 Derwent World Patents Index (DWPI) Revises Indexing

of Author Abstracts

NEWS 6 FEB 16 New FASTA Display Formats Added to USGENE and PCTGEN

NEWS 7 FEB 16 INPADOCDB and INPAFAMDB Enriched with New Content

and Features

NEWS 8 FEB 16 INSPEC Adding Its Own IPC codes and Author's E-mail

Addresses

NEWS 9 APR 02 CAS Registry Number Crossover Limits Increased to

500,000 in Key STN Databases

NEWS 10 APR 02 PATDPAFULL: Application and priority number formats

enhanced

NEWS 11 APR 02 DWPI: New display format ALLSTR available NEWS 12 APR 02 New Thesaurus Added to Derwent Databases for Smooth

Sailing through U.S. Patent Codes

NEWS 13 APR 02 EMBASE Adds Unique Records from MEDLINE, Expanding

Coverage back to 1948

NEWS 14 APR 07 CA/CAplus CLASS Display Streamlined with Removal of

Pre-IPC 8 Data Fields

NEWS 15 APR 07 50,000 World Traditional Medicine (WTM) Patents Now

Available in CAplus

NEWS 16 APR 07 MEDLINE Coverage Is Extended Back to 1947

NEWS EXPRESS FEBRUARY 15 10 CURRENT WINDOWS VERSION IS V8.4.2.

AND CURRENT DISCOVER FILE IS DATED 15 JANUARY 2010.

NEWS HOURS STN Operating Hours Plus Help Desk Availability NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN customer agreement. This agreement limits use to scientific research. Use

for software development or design, implementation of commercial

gateways, or use of CAS and STN data in the building of

products is prohibited and may result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 14:12:49 ON 20 APR 2010

=> file caplus

COST IN U.S. DOLLARS SINCE FILE

TOTAL

ENTRY SESSION

FULL ESTIMATED COST 0.22 0.22

FILE 'CAPLUS' ENTERED AT 14:13:06 ON 20 APR 2010 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS. COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLI SHER (PB) field (available

for records published or updated in Chemical Abstracts after December

26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching

databases on STN. Any dissemination, distribution, copying, or storing

of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 20 Apr 2010 VOL 152 ISS 17
FILE LAST UPDATED: 18 Apr 2010 (20100418/ED)
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Feb 2010
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE:
Feb 2010

CAplus now includes complete International Patent Classification (IPC)

reclassification data for the first quarter of 2010.

CAS Information Use Policies apply and are available at:

http://www.cas.org/legal/infopolicy.html

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s ((capture(w)(probe# or oligonucleotide# or phase)) and hybridi? and (ligate# or ligase# or ligation or lcr or ldr))/bi,ab

112836 CAPTURE/BI

94056 CAPTURE/AB

410510 PROBE#/BI

355733 PROBE#/AB 104504 OLI GONUCLEOTI DE#/BI

77279 OLI GONUCLEOTI DE# / AB

2089580 PHASE/BI

1824278 PHASE/AB

1342 CAPTURE(W)(PROBE# OR OLIGONUCLEOTIDE# OR PHASE)

214568 HYBRI DI ?/BI

180471 HYBRI DI ?/AB

18833 LI GATE#/BI

17966 LIGATE#/AB

19765 LIGASE#/BI

15407 LIGASE#/AB (FILE 'HOME' ENTERED AT 14:12:49 ON 20 APR 2010) 36532 LIGATION/BI 33697 LIGATION/AB FILE 'CAPLUS' ENTERED AT 14:13:06 ON 20 APR 2010 2846 LCR/BI L1 112 S ((CAPTURE(W)(PROBE# OR OLIGONUCLEOTI DE# OR PHASE)) AND HYBRI D 2058 LCR/AB 106 S L1 NOT 2010/PY 813 LDR/BI 12 76 S L2 NOT 2009/PY 773 LDR/AB L3 112 ((CAPTURE(W)(PROBE# OR OLIGONUCLEOTIDE# 61 S L3 NOT 2008/PY L4 OR PHASE)) AND HYBRIDI? 15 56 ST4 NOT 2007/PY AND (LIGATE# OR LIGASE# OR LIGATION OR LCR OR 46 S L5 NOT 2006/PY L6 LDR))/BI,AB 37 S L6 NOT 2005/PY L7 29 S L7 NOT 2004/PY 18 => s l1 not 2010/py23 S L8 NOT 2003/PY L9 505082 2010/PY L10 19 S L9 NOT 2002/PY 106 L1 NOT 2010/PY 13 S L10 NOT 2001/PY 111 112 10 S L11 NOT 2000/PY 9 S L12 NOT 1999/PY => s 12 not 2009/pyI 13 1845332 2009/PY 6 S L13 NOT 1998/PY 114 76 L2 NOT 2009/PY => d l14 1-6 bib ab => s |3 not 2008/py1797450 2008/PY L14 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN 61 L3 NOT 2008/PY AN 1998:138263 CAPLUS << LOGINID::20100420>> DN 128:290724 => s | 4 not 2007/pyOREF 128:57463a,57466a 1721288 2007/PY TI Rapid and sensitive detection of Chlamydia trachomatis using 56 L4 NOT 2007/PY a ligatable binary RNA probe and Q.beta. replicase AU Stefano, James E.; Genovese, Louis; An, Qi; Lu, Ling; => s 15 not 2006/py1587214 2006/PY Mccarty, Janice; Du, Yan; Stefano, Kyriaki; Burg, J. Lawrence; King, Walter; Lane, L6 46 L5 NOT 2006/PY David J => s | 6 not 2005/pvCS GENE-TRAK, Inc., Framingham, MA, 01701, USA 1433409 2005/PY SO Molecular and Cellular Probes (1997), 11(6), 407-426 37 L6 NOT 2005/PY CODEN: MCPRE6; ISSN: 0890-8508 17 PB Academic Press Ltd. => s 17 not 2004/pyDT Journal LA English 1352559 2004/PY 29 L7 NOT 2004/PY AB A simple assay format was developed for the direct detection of C. trachomatis rRNA utilizing ***ligation*** of recombinant => s l8 not 2003/py 1272622 2003/PY MDV-1 probe RNA fragments *** hybridized*** to 23S rRNA after 23 L8 NOT 2003/PY 19 capture and release => s 19 not 2002/pyfrom a solid support. Assay background (equiv. to 104 1181744 2002/PY targets) was 19 L9 NOT 2002/PY suppressed by blocking sequences in the 5' MDV reporter probe fragment => s I10 not 2001/py complementary to the 3' fragment by prehybridization of a 1124940 2001/PY 13 L10 NOT 2001/PY oligonucleotide. A pair of reporter fragments bearing a deletion within => s l11 not 2000/py the region, obtained by a hydrid-selection-amplification 1046968 2000/PY protocol, yielded 10 L11 NOT 2000/PY a low level of assay background which was reduced to <2% L12 with a blocker => s l12 not 1999/py directed against the remaining pairing sequence. This probe 965496 1999/PY set showed a 9 L12 NOT 1999/PY sensitivity of 103 mols. of 23S rRNA (>95% responding) and 113 could detect a => s l13 not 1998/py single elementary body (EB) of Chlamydia trachomatis or 1-10 935226 1998/PY EB added to a L14 6 L13 NOT 1998/PY clin. matrix of pooled neg. human cervical swab samples. The time of => d his first appearance of amplification products by real-time

fluorescence

detection showed a linear response to log increases in the target level

over a 105-fold range, permitting the detn. of target level within an

order of magnitude. The assay showed .apprx. 109-fold discrimination over

Chlamydia pneumoniae (TWAR) rRNA. High levels of cultured Candida

albicans, Escherichia coli, Staphylococcus aureus, or Neisseria gonorrhoeae had no detectable effect on assay background or the ability to

detect a single elementary body.

OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

RE.ONT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN AN 1997:37433 CAPLUS << LOGINI D:: 20100420>>

DN 126:70794

OREF 126:13605a,13608a

TI Detection of EBV early RNA (EBER-1) in parotid pleomorphic adenomas: a

novel observation utilizing ***ligation*** -dependent PCR AU Brandwein, Margaret; Li, Hongbo; Zhang, David Y.

CS Lillian and Henry M. Stratton-Hans Popper Department of Pathology,

University of New York, NY, USA

SO International Congress Series (1996), 1114(Head and Neck Cancer: Advances

in Basic Research), 401-409

CODEN: EXMDA4; ISSN: 0531-5131

PB Elsevier

DT Journal

LA English

AB Very little is known regarding the initiation and promotion of salivary

neoplasia. We utilized the recently introduced $\ensuremath{^{***}}$ ligation $\ensuremath{^{***}}$

-dependent polymerase chain reaction (LD-PCR) to detect viral RNAs. This

technique employs two ***capture*** *** probes*** for the isolation

of target RNA. A third probe contains a complementary region to the

target sequence at each end, and a generic linker region for PCR primer

binding. This probe becomes circularized upon
hybridization to

the target and forms a covalently linked circular probe by incubation with

T4 DNA ****ligase*** . The circularized probe sequence serves as a

template for Taq polymerase. A novelty of this assay is to amplify by PCR

the probe sequence rather than the target sequence. This allows for the $\,$

facile identification of RNA by PCR without the need for the reverse

transcriptase step. We studied nine cases of snap-frozen tissue from $% \left(1\right) =\left(1\right) \left(1\right) +\left(1\right) \left(1\right) \left(1\right) +\left(1\right) \left(1\right) \left($

parotid gland and six pleomorphic adenomas by LD-PCR for the presence of $% \left(1\right) =\left(1\right) \left(1\right) +\left(1\right) \left(1\right) \left(1\right) +\left(1\right) \left(1\right)$

 $\dot{\text{EBV}}$ early RNA (EBER-1). EBER-1 was identified in six of eight parotid

tissue samples and four of six pleomorphic adenomas. Although the PCR

technique does not allow for localization within tissue (viral sequences

within tumor cells vs. circulating lymphocytes), the identification of

EBER-1 in these cases does indicate that active transcription of a

latency-assocd. viral RNA is common in the parotid gland. This may have

implications on salivary tumorigenesis.

OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L14 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1995:230412 CAPLUS << LOGINI D::20100420>>

DN 122:179542

OREF 122:32745a,32748a

TI A rapid, reliable method for detection of known point mutations:

point-EXACCT

AU Somers, Veerle A. M. C.; Moerkerk, Peter T. M.; Murtagh, James J., Jr.;

Thunnissen, Frederik B. J. M.

CS Department Pathology, University Limburg, Maastricht, Neth.

SO Nucleic Acids Research (1994), 22(22), 4840-1 CODEN: NARHAD; ISSN: 0305-1048

PB Oxford University Press

DT Journal

LA English

AB Point mutations in the human genome play a central role in tumorigenesis.

Several methods are available for detection of known point mutations. The

detection format is based on an extension of the EXACCT procedure. In

short: after exonuclease digestion, polymerase chain reaction fragments

are detd. by ***hybridization*** with a capture and a detection probe

complementary to sequences near the 3' end of the antisense fragment. The

capture *** probe*** bears a biotin residue and the other probe digoxigenin. After *** hybridization*** the PCR product

hybrids are

captured in streptavidin-coated microtiter plates and detected with

labeled anti-digoxigenin antibody. For the detection of known point

mutations this procedure was extended by using after the capture step the

*** ligation*** of a mutation-specific *** capture***

with adjacent detection probe (Point-EXACCT). Point-EXACCT requires

considerably less time and effort than other techniques used for the

detection of known point mutations. This method is easily automated

permitting rapid screening of tissue banks with multiple probes to

individual base substitutions, deletions or addns. The simplicity of

Point-EXACCT makes it a highly promising method for the detection of known

point mutations. AN 1994:70885 CAPLUS < LOGINID::20100420>> OSC.G 15 THERE ARE 15 CAPLUS RECORDS THAT CITE THIS DN 120:70885 OREF 120:12639a,12642a RECORD (15 CITINGS) TI Chlamydiae probes for use in solution phase sandwich L14 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN * * * hybridization * * * AN 1994:70887 CAPLUS << LOGINI D::20100420>> assavs DN 120:70887 IN Sanchez-Pescador, Ray; Besemer, Diana J.; Urdea, Michael OREF 120:12639a,12642a S. TI Cytomegalovirus (CMV) probes for use in solution phase PA Chiron Corp., USA sandwich SO PCT Int. Appl., 84 pp. * * * hybridization* * * assays CODEN: PLXXD2 IN Kolberg, Janice A.; Shen, Lu Ping; Urdea, Michael S. DT Patent PA Chiron Corp., USA LA English SO PCT Int. Appl., 71 pp. FAN. CNT 1 CODEN: PIXXD2 KIND DATE APPLICATION NO. PATENT NO. DT Patent LA English -----.... FAN. CNT 1 PI WO 9313221 A1 19930708 WO 1992-US11035 KIND DATE APPLICATION NO. PATENT NO. 19921222 W: AU, CA, JP, KR DATE RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, PI WO 9313227 A1 19930708 WO 1992-US11170 NL, PT, SE AU 9334672 19921222 19930728 AU 1993-34672 W: CA, JP, KR 19921222 RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, EP 726963 A1 19960821 EP 1993-903387 NL. PT. SE 19921222 A1 19941123 EP 1993-902723 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, EP 625214 NL, PT, SE R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, US 5618674 19970408 US 1995-479487 NL, PT, SE 19950607 US 5407795 A 19950418 US 1993-138608 PRAI US 1991-813587 A 19911223 A 19921222 WO 1992-US11035 19931015 ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS PRAI US 1991-813590 A 19911223 WO 1992-US11170 W 19921222 DISPLAY FORMAT ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS AB The title probes, i.e. amplifier probe and ***capture*** DISPLAY FORMAT * * * probe* * * AB The title probes, i.e. amplifier or ***capture*** , comprise a first segment with nucleotide sequence **probes*** substantially comprises a nucleotide sequence complementary to a complementary to a segment of Chlamydiae plasmid DNA and segment of CMV nucleic a second segment acid and a nucleotide sequence complementary to a segment with nucleotide sequence substantially complementary to an of nucleotide oliaonucleotide sequence of an amplifier multimer or a capture solid phase, multimer or an oligonucleotide bound to a solid phase, resp. Thus, a resp. Thus, a comb-type polynucleotide having 15 branch sites and side comb-type polynucleotide having 15 branch sites and side chain extensions chain extensions having 3 labeled probe binding sites was prepd. as an having 3 labeled probe binding sites was synthesized and amplifier multimer. used as a labeled multimer. The amplifier and ***capture*** CMV amplifier and ***capture*** *** probes*** probes*** are (contg., in addn. to *** hybridized*** with sample, the formed complexes are sequences complementary to CMV sequences, a 5' extension captured by complementary to the amplifier multimer or a downstream sequence of oligonucleotide-bound solid phase, and the captured CTTCTTTGGAGAAAGTGGTG complexes are *** hybridized*** with the oligonucleotide multimer and complementary to an immobilized oligonucleotide, resp.) were complementary used along with the amplifier multimer and capture solid phase in a labeled oligonucleotide for Chlamydiae detection. THERE ARE 7 CAPLUS RECORDS THAT CITE THIS OSC.G 7 * * * hybridization* * * assay of CMV. RECORD (7 CITINGS) OSC.G 10 THERE ARE 10 CAPLUS RECORDS THAT CITE THIS RE ONT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR RECORD (10 CITINGS) THIS RECORD RE.ONT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR ALL CITATIONS AVAILABLE IN THE RE FORMAT THIS RECORD L14 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1993:663642 CAPLUS << LOGINI D:: 20100420>>

DN 119:263642

OREF 119:46973a.46976a

TI A transcriptionally amplified DNA probe assay with ligatable probes and

immunochemical detection

AU Carpenter, William R.; Schutzbank, Ted E.; Tevere, Vincent J.; Tocyloski,

Kenneth R.; Dattagupta, Nanibushan; Yeung, Kwok K.

CS Diagn. Div., Miles Inc., Tarrytown, NY, 10591, USA

SO Clinical Chemistry (Washington, DC, United States) (1993), 39(9), 1934-8

CODEN: CLCHAU; ISSN: 0009-9147

DT Journal

LA English

AB Transcriptionally amplified DNA probes are valuable tools in the

development of sensitive nucleic acid-based diagnostic assays. Here the

authors describe a model assay using a novel oligonucleotide hairpin probe

that encodes a T7 RNA polymerase promoter. The hairpin probe and an $\,$

captured onto streptavidin-coated magnetic particles. After

*** ligation*** of the immobilized probes, which served to
aintain

specificity, the hairpin probe was transcribed by T7 RNA polymerase. The $\,$

amplified RNA product was *** hybridized*** to the
capture

*** probe*** and bound to the streptavidin-coated magnetic particles.

The immobilized heteroduplex was detected with an antibody-alk.

phosphatase conjugate specific for DNA: RNA hybrids, and the chemiluminescent substrate adamantyl-1,2-dioxetane Ph phosphate. Ten

attomoles of target DNA could be detected in a background of 5 .mu.g of

unrelated DNA. The chemiluminescent immunoassay was as sensitive as

radioactive detection of specific product after gel electrophoresis.

OSC.G. 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

=> log y

COST ÎN U.S. DOLLARS SINCE FILE TOTAL

ENTRY SESSION

FULL ESTIMATED COST 97.83 98.05

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS) SINCE

FILE TOTAL

ENTRY SESSION

CA SUBSCRIBER PRICE -5.10 -5.10

STN INTERNATIONAL LOGOFF AT 14:16:35 ON 20 APR 2010